.-online.com antibodies

Datasheet for ABIN7145737 anti-BSX antibody (AA 1-233)

Image



Overview

Quantity:	100 µg
Target:	BSX
Binding Specificity:	AA 1-233
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BSX antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Brain-specific homeobox protein homolog protein (1-233AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified


Target Details

Target:	BSX
Alternative Name:	BSX (BSX Products)
Background:	Background: DNA binding protein that function as transcriptional activator. Is essentiel for
	normal postnatal growth and nursing. Is an essential factor for neuronal neuropeptide Y and

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7145737 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
	agouti-related peptide function and locomotory behavior in the control of energy balance (By similarity). Aliases: BSX antibody, BSX1Brain-specific homeobox protein homolog antibody
UniProt:	Q3C1V8
Pathways:	Feeding Behaviour
Application Details	
Application Notes:	Recommended dilution: IHC:1:500-1:1000,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7145737 diluted at 1:500 and staining in paraffin-embedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.