# antibodies .- online.com





### Datasheet for ABIN7145765

# anti-BARD1 antibody (AA 428-777) (HRP)



#### Overview

Quantity:	100 μg
Target:	BARD1
Binding Specificity:	AA 428-777
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BARD1 antibody is conjugated to HRP
Application:	ELISA

#### **Product Details**

Immunogen:	Recombinant Human BRCA1-associated RING domain protein 1 protein (428-777AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	BARD1
Alternative Name:	BARD1 (BARD1 Products)
Background:	Background: E3 ubiquitin-protein ligase. The BRCA1-BARD1 heterodimer specifically mediates
	the formation of \'Lys-6\'-linked polyubiquitin chains and coordinates a diverse range of cellular

pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability. Plays a central role in the control of the cell cycle in response to DNA damage. Acts by mediating ubiquitin E3 ligase activity that is required for its tumor suppressor function. Also forms a heterodimer with CSTF1/CSTF-50 to modulate mRNA processing and RNAP II stability by inhibiting pre-mRNA 3\' cleavage.

Aliases: BARD-1 antibody, Bard1 antibody, BARD1\_HUMAN antibody, BRCA1 associated RING domain 1 antibody, BRCA1 associated RING domain gene 1 antibody, BRCA1 associated RING domain protein 1 antibody, BRCA1-associated RING domain protein 1 antibody

UniProt: Q99728

Pathways: DNA Damage Repair

#### **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.