# antibodies - online.com







# anti-CCL14 antibody (AA 20-93)



Image



#### Overview

Quantity:	100 μg
Target:	CCL14
Binding Specificity:	AA 20-93
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCL14 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

### **Product Details**

Immunogen:	Recombinant Human C-C motif chemokine 14 protein (20-93AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	CCL14
Alternative Name:	CCL14 (CCL14 Products)
Background:	Background: Has weak activities on human monocytes and acts via receptors that also
	recognize MIP-1 alpha. It induced intracellular Ca2+ changes and enzyme release, but no

neutrophils, and eosinophil leukocytes. Enhances the proliferation of CD34 myeloid progenitor cells. The processed form HCC-1(9-74) is a chemotactic factor that attracts monocytes eosinophils, and T-cells and is a ligand for CCR1, CCR3 and CCR5.

Aliases: CC-1 antibody, CC-3 antibody, CCL14 antibody, CCL14\_HUMAN antibody, chemokine (C-C motif) ligand 14 antibody, Chemokine CC-1/CC-3 antibody, chemokine CC1 antibody, chemokine CC3 antibody, chemokine HCC1 antibody, chemokine HCC3 antibody, CKb1 antibody, HCC 1 antibody, HCC 3 antibody, HCC-1(1-74) antibody, HCC-1(9-74) antibody, HCC-1/HCC-3 antibody, HCC-3 antibody, HCC1 antibody, HEMOFILTRATE CC CHEMOKINE 1 antibody, MCIF antibody, NCC-2 antibody, NCC2 antibody, new CC chemokine 2 antibody, SCYA14 antibody, SCYL2 antibody, small inducible cytokine subfamily A (Cys-Cys), member 14

chemotaxis, at concentrations of 100-1,000 nM, and was inactive on T-lymphocytes,

UniProt:

Q16627

#### Application Details

Application Notes:	Recommended dilution: IHC:1:20-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid

antibody, Small-inducible cytokine A14 antibody, SY14 antibody


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



#### **Immunohistochemistry**

**Image 1.** Immunohistochemistry of paraffin-embedded human kidney tissue using ABIN7145954 at dilution of 1:100