



[Go to Product page](#)

Datasheet for ABIN7145983
anti-CCL22 antibody (AA 25-82)

1 Image

Overview

Quantity:	100 µg
Target:	CCL22
Binding Specificity:	AA 25-82
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCL22 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Recombinant Human C-C motif chemokine 22 protein (25-82AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CCL22
Alternative Name:	CCL22 (CCL22 Products)
Background:	Background: May play a role in the trafficking of activated/effector T-lymphocytes to inflammatory sites and other aspects of activated T-lymphocyte physiology. Chemotactic for

Target Details

monocytes, dendritic cells and natural killer cells. Mild chemoattractant for primary activated T-lymphocytes and a potent chemoattractant for chronically activated T-lymphocytes but has no chemoattractant activity for neutrophils, eosinophils, and resting T-lymphocytes. Binds to CCR4. Processed forms MDC(3-69), MDC(5-69) and MDC(7-69) seem not be active.

Aliases: A 152E5.1 antibody, ABCD 1 antibody, ABCD1 antibody, C C motif chemokine 22 antibody, CC chemokine STCP 1 antibody, CC chemokine STCP-1 antibody, ccl 22 antibody, Ccl22 antibody, CCL22_HUMAN antibody, Chemokine (C C motif) ligand 22 antibody, DC/B CK antibody, DCBCK antibody, Macrophage-derived chemokine antibody, MDC antibody, MDC(1-69) antibody, MDC(7-69) antibody, MGC34554 antibody, SCYA22 antibody, Small inducible cytokine subfamily A (Cys Cys) member 22 antibody, Small inducible cytokine subfamily A, member 22 antibody, Small-inducible cytokine A22 antibody, STCP 1 antibody, STCP1 antibody, Stimulated T cell chemotactic protein 1 antibody, Stimulated T-cell chemotactic protein 1 antibody

UniProt: [000626](#)

Application Details

Application Notes: Recommended dilution: WB:1:500-1:5000,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Western Blotting

Image 1. Western blot All lanes: CCL22 antibody at 2 μ g/mL Lane 1: 293T whole cell lysate Lane 2: EC109 whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/15000 dilution Predicted band size: 11 kDa Observed band size: 50, 80 kDa