# antibodies .- online.com





## anti-CCL22 antibody (AA 25-82) (HRP)



#### Overview

Quantity:	100 μg
Target:	CCL22
Binding Specificity:	AA 25-82
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCL22 antibody is conjugated to HRP
Application:	ELISA

#### **Product Details**

Immunogen:	Recombinant Human C-C motif chemokine 22 protein (25-82AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	CCL22
Alternative Name:	CCL22 (CCL22 Products)
Background:	Background: May play a role in the trafficking of activated/effector T-lymphocytes to
	inflammatory sites and other aspects of activated T-lymphocyte physiology. Chemotactic for

monocytes, dendritic cells and natural killer cells. Mild chemoattractant for primary activated T-lymphocytes and a potent chemoattractant for chronically activated T-lymphocytes but has no chemoattractant activity for neutrophils, eosinophils, and resting T-lymphocytes. Binds to CCR4. Processed forms MDC(3-69), MDC(5-69) and MDC(7-69) seem not be active.

Aliases: A 152E5.1 antibody, ABCD 1 antibody, ABCD1 antibody, C C motif chemokine 22 antibody, CC chemokine STCP 1 antibody, CC chemokine STCP-1 antibody, ccl 22 antibody, Ccl22 antibody, CCL22\_HUMAN antibody, Chemokine (C C motif) ligand 22 antibody, DC/B CK antibody, DCBCK antibody, Macrophage-derived chemokine antibody, MDC antibody, MDC(1-69) antibody, MDC(7-69) antibody, MGC34554 antibody, SCYA22 antibody, Small inducible cytokine subfamily A (Cys Cys) member 22 antibody, Small inducible cytokine subfamily A, member 22 antibody, Small-inducible cytokine A22 antibody, STCP 1 antibody, STCP1 antibody, Stimulated T cell chemotactic protein 1 antibody

UniProt:

000626

#### **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.