



[Go to Product page](#)

Datasheet for ABIN7145992
anti-CCL27 antibody (AA 25-112)

1 Image

Overview

Quantity:	100 µg
Target:	CCL27
Binding Specificity:	AA 25-112
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCL27 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human C-C motif chemokine 27 protein (25-112AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CCL27
Alternative Name:	CCL27 (CCL27 Products)
Background:	Background: Chemotactic factor that attracts skin-associated memory T-lymphocytes. May play a role in mediating homing of lymphocytes to cutaneous sites. Binds to CCR10.

Target Details

Aliases: ALP antibody, C-C motif chemokine 27 antibody, CC chemokine ILC antibody, Ccl27 antibody, CCL27_HUMAN antibody, Chemokine (C-C motif) ligand 27 antibody, CTACK antibody, CTAK antibody, Cutaneous T cell attracting chemokine antibody, Cutaneous T-cell-attracting chemokine antibody, ESkin antibody, IL 11 R alpha locus chemokine antibody, IL-11 R-alpha-locus chemokine antibody, ILC antibody, PESKY antibody, SCYA27 antibody, Skinkine antibody, Small inducible cytokine A27 antibody, small inducible cytokine subfamily A (Cys-Cys), member 27 antibody, Small-inducible cytokine A27 antibody

UniProt: [Q9Y4X3](#)

Application Details

Application Notes: Recommended dilution: IHC:1:20-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7145992 diluted at 1:100 and staining in paraffin-embedded human lymph node tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.