antibodies - online.com







anti-CELSR3 antibody (AA 514-718)



Image



Overview

Quantity:	100 μL
Target:	CELSR3
Binding Specificity:	AA 514-718
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CELSR3 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Cadherin EGF LAG seven-pass G-type receptor 3 protein (514-718AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CELSR3
Alternative Name:	CELSR3 (CELSR3 Products)
Background:	Background: Receptor that may have an important role in cell/cell signaling during nervous system formation.

Target Details

Aliases: Anchor protein antibody, Cadherin EGF LAG seven-pass G-type receptor 3 antibody, Cadherin family member 11 antibody, CDHF11 antibody, CELR3_HUMAN antibody, Celsr3 antibody, EGF like domain multiple 1 antibody, EGF-like protein 1 antibody, EGFL1 antibody, Egidermal growth factor-like protein 1 antibody, Flamingo antibody, Flamingo homolog 1 antibody, FMI1 antibody, hFmi1 antibody, MEGF2 antibody, Multiple EGF-like domains protein 2 antibody, Multiple epidermal growth factor-like domains protein 2 antibody, RESDA1 antibody

UniProt:

Q9NYQ7

Application Details

1010 - 200 -		
Application Notes:	Recommended dilution: IHC:1:200-1:500,	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	

Buffer: Preservative: 0.03 % Proclin 300


Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7146186 diluted at 1:300 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.