.-online.com antibodies

Datasheet for ABIN7146604 anti-PRKACB antibody (Catalytic Subunit beta) (FITC)



Overview

Quantity:	100 µg
Target:	PRKACB
Binding Specificity:	AA 2-53, Catalytic Subunit beta
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRKACB antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human cAMP-dependent protein kinase catalytic subunit beta protein (2-53AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	PRKACB
Alternative Name:	PRKACB (PRKACB Products)
Background:	Background: Mediates cAMP-dependent signaling triggered by receptor binding to GPCRs. PKA
	activation regulates diverse cellular processes such as cell proliferation, the cell cycle,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7146604 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Preservative:

Storage:

Precaution of Use:

Storage Comment:

ProClin

-20 °C,-80 °C

handled by trained staff only.

	differentiation and regulation of microtubule dynamics, chromatin condensation and
	decondensation, nuclear envelope disassembly and reassembly, as well as regulation of
	intracellular transport mechanisms and ion flux. Regulates the abundance of
	compartmentalized pools of its regulatory subunits through phosphorylation of PJA2 which
	binds and ubiquitinates these subunits, leading to their subsequent proteolysis.
	Aliases: cAMP-dependent protein kinase catalytic beta subunit isoform 4ab antibody, cAMP-
	dependent protein kinase catalytic subunit beta antibody, KAPCB_HUMAN antibody, PKA C beta
	antibody, PKA C-beta antibody, PKACB antibody, Prkacb antibody, protein kinase A catalytic
	subunit beta antibody, Protein kinase cAMP dependent catalytic beta antibody
UniProt:	P22694
Pathways:	AMPK Signaling, Hedgehog Signaling, EGFR Signaling Pathway, Neurotrophin Signaling
	Pathway, Thyroid Hormone Synthesis, Myometrial Relaxation and Contraction, M Phase, G-
	protein mediated Events, Interaction of EGFR with phospholipase C-gamma, Lipid Metabolism
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be