

Datasheet for ABIN7146860

## anti-Casein Kinase 1 delta antibody (Isoform delta) (FITC)



[Go to Product page](#)

### Overview

Quantity:	100 µg
Target:	Casein Kinase 1 delta (CSNK1D)
Binding Specificity:	AA 152-398, Isoform delta
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Casein Kinase 1 delta antibody is conjugated to FITC
Application:	Please inquire

### Product Details

Immunogen:	Recombinant Human Casein kinase I isoform delta protein (152-398AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	Casein Kinase 1 delta (CSNK1D)
Alternative Name:	CSNK1D ( <a href="#">CSNK1D Products</a> )
Background:	Background: Essential serine/threonine-protein kinase that regulates diverse cellular growth and survival processes including Wnt signaling, DNA repair and circadian rhythms. It can

## Target Details

phosphorylate a large number of proteins. Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates. Phosphorylates connexin-43/GJA1, MAP1A, SNAPIN, MAPT/TAU, TOP2A, DCK, HIF1A, EIF6, p53/TP53, DVL2, DVL3, ESR1, AIB1/NCOA3, DNMT1, PKD2, YAP1, PER1 and PER2. Central component of the circadian clock. In balance with PP1, determines the circadian period length through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. Controls PER1 and PER2 nuclear transport and degradation. YAP1 phosphorylation promotes its SCF(beta-TRCP) E3 ubiquitin ligase-mediated ubiquitination and subsequent degradation. DNMT1 phosphorylation reduces its DNA-binding activity. Phosphorylation of ESR1 and AIB1/NCOA3 stimulates their activity and coactivation. Phosphorylation of DVL2 and DVL3 regulates WNT3A signaling pathway that controls neurite outgrowth. EIF6 phosphorylation promotes its nuclear export. Triggers down-regulation of dopamine receptors in the forebrain. Activates DCK in vitro by phosphorylation. TOP2A phosphorylation favors DNA cleavable complex formation. May regulate the formation of the mitotic spindle apparatus in extravillous trophoblast. Modulates connexin-43/GJA1 gap junction assembly by phosphorylation. Probably involved in lymphocyte physiology. Regulates fast synaptic transmission mediated by glutamate.

Aliases: Casein Kinase 1 delta antibody, Casein kinase I delta isoform antibody, Casein kinase I isoform delta antibody, CKI delta antibody, CKI-delta antibody, CKId antibody, CKIdelta antibody, CSNK 1D antibody, CSNK1D antibody, HCKID antibody, KC1D\_HUMAN antibody, Protein kinase CK1 delta antibody

UniProt: [P48730](#)

Pathways: [Hedgehog Signaling, M Phase](#)

## Application Details

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

---

Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.