

Datasheet for ABIN7146891 anti-CSNK2A2 antibody (AA 8-350) (HRP)



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μg
Target:	CSNK2A2
Binding Specificity:	AA 8-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CSNK2A2 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Casein kinase II subunit alpha' protein (8-350AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CSNK2A2
Alternative Name:	CSNK2A2 (CSNK2A2 Products)
Background:	Background: Catalytic subunit of a constitutively active serine/threonine-protein kinase complex
	that phosphorylates a large number of substrates containing acidic residues C-terminal to the

phosphorylated serine or threonine. Regulates numerous cellular processes, such as cell cycle progression, apoptosis and transcription, as well as viral infection. May act as a regulatory node which integrates and coordinates numerous signals leading to an appropriate cellular response. During mitosis, functions as a component of the p53/TP53-dependent spindle assembly checkpoint (SAC) that maintains cyclin-B-CDK1 activity and G2 arrest in response to spindle damage. Also required for p53/TP53-mediated apoptosis, phosphorylating 'Ser-392' of p53/TP53 following UV irradiation. Can also negatively regulate apoptosis. Phosphorylates the caspases CASP9 and CASP2 and the apoptotic regulator NOL3. Phosphorylation protects CASP9 from cleavage and activation by CASP8, and inhibits the dimerization of CASP2 and activation of CASP8. Regulates transcription by direct phosphorylation of RNA polymerases I, II, III and IV. Also phosphorylates and regulates numerous transcription factors including NFkappa-B, STAT1, CREB1, IRF1, IRF2, ATF1, SRF, MAX, JUN, FOS, MYC and MYB. Phosphorylates Hsp90 and its co-chaperones FKBP4 and CDC37, which is essential for chaperone function. Regulates Wnt signaling by phosphorylating CTNNB1 and the transcription factor LEF1. Acts as an ectokinase that phosphorylates several extracellular proteins. During viral infection, phosphorylates various proteins involved in the viral life cycles of EBV, HSV, HBV, HCV, HIV, CMV and HPV.

Aliases: Casein kinase 2 alpha prime polypeptide antibody, Casein kinase II alpha' chain antibody, Casein kinase II subunit alpha' antibody, CK II alpha' antibody, CK II antibody, CK2A2 antibody, CKII antibody, CSK22_HUMAN antibody, CSNK2A1 antibody, CSNK2A2 antibody, FLJ43934 antibody

UniProt: P19784

Pathways: SARS-CoV-2 Protein Interactome

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format:

Buffer:

Preservative: 0.03 % Proclin 300

Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.