antibodies - online.com







anti-FLIP antibody (AA 1-250)





()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Quantity:	100 μg
Target:	FLIP (CFLAR)
Binding Specificity:	AA 1-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FLIP antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human CASP8 and FADD-like apoptosis regulator protein (1-250AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	FLIP (CFLAR)
Alternative Name:	CFLAR (CFLAR Products)
Background:	Background: Apoptosis regulator protein which may function as a crucial link between cell survival and cell death pathways in mammalian cells. Acts as an inhibitor of TNFRSF6

signaling complex (DISC) thereby blocking further recruitment and processing of caspase-8 at the complex. Full length and shorter isoforms have been shown either to induce apoptosis or to reduce TNFRSF-triggered apoptosis. Lacks enzymatic (caspase) activity. Aliases: c FLIP antibody, c FLIPL antibody, c FLIPS antibody, c FLIPS antibody, c-FLIPL antibody, c-FLIPR antibody, c-FLIPS antibody, CASH antibody, CASP8 and FADD like apoptosis regulator antibody, CASP8 and FADD-like apoptosis regulator subunit p12 antibody, CASP8AP1 antibody, Caspase homolog antibody, Caspase like apoptosis regulatory protein antibody, Caspase related inducer of apoptosis antibody, Caspase-eight-related protein antibody, Caspase-like apoptosis regulatory protein antibody, Casper antibody, cellular flice inhibitory protein antibody, Cellular FLICE-like inhibitory protein antibody, CFLA antibody, Cflar antibody, CFLAR_HUMAN antibody, CLARP antibody, FADD like anti apoptotic molecule antibody, FADD-like antiapoptotic molecule 1 antibody, FLAME antibody, FLAME-1 antibody, FLAME1 antibody, FLIP antibody, I FLICE antibody, I-FLICE antibody, Inhibitor of FLICE antibody, MACH-related inducer of toxicity antibody, MRIT antibody, OTTHUMP00000163715 antibody, OTTHUMP00000206475 antibody, OTTHUMP00000206476 antibody, OTTHUMP00000206478 antibody, OTTHUMP00000206479 antibody, OTTHUMP00000206480 antibody, OTTHUMP00000206482 antibody, OTTHUMP00000207360 antibody, Usurpin antibody, Usurpin beta antibody

mediated apoptosis. A proteolytic fragment (p43) is likely retained in the death-inducing

LluiDuit.	015510
UniProt:	015519

Pathways: Apoptosis, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development

Application Details


Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500, IF:1:20-1:200,
Restrictions:	For Research Use only
Handling	

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:	-20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunofluorescence

Image 1. Immunofluorescence staining of Hela cells with ABIN7146901 at 1:40, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Western Blotting

Image 2. Western Blot Positive WB detected in: Jurkat whole cell lysate, K562 whole cell lysate, 293T whole cell lysate All lanes: CFLAR antibody at 4 μg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 56, 26, 42, 28, 52, 51, 40, 31, 35, 53, 34, 24, 50, 45 kDa Observed band size: 56 kDa



Immunohistochemistry

Image 3. IHC image of ABIN7146901 diluted at 1:400 and staining in paraffin-embedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

35	
	Please check the product details page for more images. Overall 4 images are available for ABIN7146901.