



[Go to Product page](#)

Datasheet for ABIN7147090

anti-CNOT1 antibody (AA 604-867) (Biotin)

Overview

Quantity:	100 µg
Target:	CNOT1
Binding Specificity:	AA 604-867
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CNOT1 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human CCR4-NOT transcription complex subunit 1 protein (604-867AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CNOT1
Alternative Name:	CNOT1 (CNOT1 Products)
Background:	Background: Scaffolding component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA

Target Details

degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation. Additional complex functions may be a consequence of its influence on mRNA expression. Its scaffolding function implies its interaction with the catalytic complex module and diverse RNA-binding proteins mediating the complex recruitment to selected mRNA 3'UTRs. Involved in degradation of AU-rich element (ARE)-containing mRNAs probably via association with ZFP36. Mediates the recruitment of the CCR4-NOT complex to miRNA targets and to the RISC complex via association with TNRC6A, TNRC6B or TNRC6C. Acts as a transcriptional repressor. Represses the ligand-dependent transcriptional activation by nuclear receptors. Involved in the maintenance of embryonic stem (ES) cell identity. Aliases: CNOT1 antibody, CDC39 antibody, KIAA1007 antibody, NOT1 antibody, AD-005CCR4-NOT transcription complex subunit 1 antibody, CCR4-associated factor 1 antibody, Negative regulator of transcription subunit 1 homolog antibody, NOT1H antibody, hNOT1 antibody

UniProt: [A5YKK6](#)

Pathways: [Retinoic Acid Receptor Signaling Pathway](#), [Intracellular Steroid Hormone Receptor Signaling Pathway](#), [Regulation of Intracellular Steroid Hormone Receptor Signaling](#), [Nuclear Hormone Receptor Binding](#), [Stem Cell Maintenance](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.