## antibodies .- online.com





## anti-ARHGAP24 antibody (AA 501-600) (Cy7)



Go to Product page

( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL
Target:	ARHGAP24
Binding Specificity:	AA 501-600
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ARHGAP24 antibody is conjugated to Cy7
Application:	Western Blotting (WB)

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ARHGAP24	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat	
Purification:	Purified by Protein A.	

## Target Details

Target:	ARHGAP24	
Alternative Name:	ARHGAP24 (ARHGAP24 Products)	
Background:	Synonyms: Rho GTPase-activating protein 24, Filamin-A-associated RhoGAP, FilGAP, RAC1-	
	and CDC42-specific GTPase-activating protein of 72 kDa, RC-GAP72, Rho-type GTPase-	

	activating protein 24, RhoGAP of 73 kDa, Sarcoma antigen NY-SAR-88, p73RhoGAP,		
	ARHGAP24, FILGAP		
	Background: Rho GTPase-activating protein involved in cell polarity, cell morphology and		
	cytoskeletal organization. Acts as a GTPase activator for the Rac-type GTPase by converting it		
	to an inactive GDP-bound state. Controls actin remodeling by inactivating Rac downstream of		
	Rho leading to suppress leading edge protrusion and promotes cell retraction to achieve		
	cellular polarity. Able to suppress RAC1 and CDC42 activity in vitro. Overexpression induces cell		
	rounding with partial or complete disruption of actin stress fibers and formation of membrane		
	ruffles, lamellipodia, and filopodia. Isoform 2 is a vascular cell-specific GAP involved in		
	modulation of angiogenesis.		
Gene ID:	83478		
UniProt:	Q8N264		
Pathways:	Regulation of Cell Size, Positive Regulation of Response to DNA Damage Stimulus		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Expiry Date:	12 months		