antibodies -online.com





anti-CDCA7L antibody (AA 14-167) (FITC)



Go to Product page

\sim				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μg
Target:	CDCA7L
Binding Specificity:	AA 14-167
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDCA7L antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Cell division cycle-associated 7-like protein (14-167AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	CDCA7L	
Alternative Name:	CDCA7L (CDCA7L Products)	
Background:	Background: Plays a role in transcriptional regulation as a repressor that inhibits monoamine	
	oxidase A (MAOA) activity and gene expression by binding to the promoter. Plays an important	

Target Details

oncogenic role in mediating the full transforming effect of MYC in medulloblastoma cells. Involved in apoptotic signaling pathways, May act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1.

Aliases: c Myc Interacting protein JPO2 antibody, CDA7L_HUMAN antibody, Cdca7l antibody, Cell division cycle-associated 7-like protein antibody, DKFZp762L0311 antibody, HR1 antibody, JPO2 antibody, Protein JPO2 antibody, R1 antibody, RAM2 antibody, Transcription factor RAM2

UniProt:

Q96GN5

antibody

Application Details

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	