# antibodies .- online.com







## anti-CHMP6 antibody (AA 2-201) (FITC)



#### Overview

| Quantity:            | 100 μg                                    |
|----------------------|---|
| Target:              | CHMP6                                     |
| Binding Specificity: | AA 2-201                                  |
| Reactivity:          | Human                                     |
| Host:                | Rabbit                                    |
| Clonality:           | Polyclonal                                |
| Conjugate:           | This CHMP6 antibody is conjugated to FITC |
| Application:         | Please inquire                            |

#### **Product Details**

| Immunogen:        | Recombinant Human Charged multivesicular body protein 6 protein (2-201AA) |
|-------------------|---|
| Isotype:          | IgG   |
| Cross-Reactivity: | Human   |
| Purification:     | >95%, Protein G purified  |

### Target Details

| Target:           | CHMP6   |
|-------------------|---|
| Alternative Name: | CHMP6 (CHMP6 Products)  |
| Background:       | Background: Probable core component of the endosomal sorting required for transport             |
|                   | complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting |

of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-0, -I,-II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and the budding of enveloped viruses (HIV-1 and other lentiviruses). ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. In the ESCRT-III complex, it probably serves as an acceptor for the ESCRT-II complex on endosomal membranes. Aliases: Charged multivesicular body protein 6 antibody, chmp6 antibody, CHMP6\_HUMAN antibody, Chromatin modifying protein 6 antibody, Chromatin-modifying protein 6 antibody, FLJ11749 antibody, hVps 20 antibody, hVps20 antibody, novel SNF7 protein antibody, OTTMUSP00000004363 antibody, RGD1565325 antibody, RP24 225K18.4 antibody, Vacuolar protein sorting associated protein 20 antibody, Vacuolar protein sorting-associated protein 20 antibody, Vps 20 antibody, Vps 20 antibody

UniProt:

Q96FZ7

#### **Application Details**

Restrictions:

For Research Use only

#### Handling

| Format:            | Liquid  |
|--------------------|---|
| Buffer:            | Preservative: 0.03 % Proclin 300<br>Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4                                |
| Preservative:      | ProClin   |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | -20 °C,-80 °C   |
| Storage Comment:   | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |