antibodies.com

Datasheet for ABIN7147545 anti-HUS1B antibody (AA 37-113)

Image



Overview

Quantity:	100 μL
Target:	HUS1B
Binding Specificity:	AA 37-113
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HUS1B antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Checkpoint protein HUS1B protein (37-113AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	HUS1B
Alternative Name:	HUS1B (HUS1B Products)
Background:	Background: checkpoint clamp complex, site of double-strand break, double-strand break repair
	via homologous recombination, intra-S DNA damage checkpoint, meiotic DNA integrity

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7147545 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details		
	checkpoint, mitotic DNA replication checkpoint, nucleotide-excision repair Aliases: HUS1B antibody, Checkpoint protein HUS1B antibody, hHUS1B antibody	
UniProt:	Q8NHY5	
Application Details		
Application Notes:	Recommended dilution: IHC:1:200-1:500,	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	

Images



Immunohistochemistry

Image 1. IHC image of ABIN7147545 diluted at 1:400 and staining in paraffin-embedded human prostate tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.