antibodies -online.com





anti-CHD5 antibody (AA 1600-1699)

3 Images



Go to Product page

\sim					
	1//	⊃r	V/I	Φ\	Λ

Quantity:	100 μg	
Target:	CHD5	
Binding Specificity:	AA 1600-1699	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CHD5 antibody is un-conjugated	
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)	
Product Details		
Immunogen:	Recombinant Human Chromodomain-helicase-DNA-binding protein 5 protein (1600-1699AA)	

Immunogen:	Recombinant Human Chromodomain-helicase-DNA-binding protein 5 protein (1600-1699AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CHD5
Alternative Name:	CHD5 (CHD5 Products)
Background:	Background: Chromatin-remodeling protein that binds DNA through histones and regulates
	gene transcription. May specifically recognize and bind trimethylated \'Lys-27\' (H3K27me3)

and non-methylated \'Lys-4\' of histone H3. Plays a role in the development of the nervous system by activating the expression of genes promoting neuron terminal differentiation. In parallel, it may also positively regulate the trimethylation of histone H3 at \'Lys-27\' thereby specifically repressing genes that promote the differentiation into non-neuronal cell lineages. Tumor suppressor, it regulates the expression of genes involved in cell proliferation and differentiation. Downstream activated genes may include CDKN2A that positively regulates the p53/TP53 pathway, which in turn, prevents cell proliferation. In spermatogenesis, it probably regulates histone hyperacetylation and the replacement of histones by transition proteins in chromatin, a crucial step in the condensation of spermatid chromatin and the production of functional spermatozoa.


Aliases: ATP-dependent helicase CHD5 antibody, CHD 5 antibody, CHD-5 antibody, CHD5 antibody, CHD5_HUMAN antibody, Chromodomain-helicase-DNA-binding protein 5 antibody, DKFZp434N231 antibody, KIAA0444 antibody, RP1-233K16.2 antibody

UniProt:

Q8TDI0


Application Details

Application Notes:	Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunofluorescence

Image 1. Immunofluorescent analysis of Hela cells using ABIN7147724 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Immunohistochemistry

Image 2. Immunohistochemistry of paraffin-embedded human liver cancer using ABIN7147724 at dilution of 1:100



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human placenta tissue using ABIN7147724 at dilution of 1:100