

Datasheet for ABIN7147842

anti-BHLHE41 antibody (AA 185-285) (Biotin)



Go to Product page

| _ | | | | |
|-----|-----|-----|---|-----|
| () | VE | r\/ | | Λ/ |
| () | V C | I V | 1 | v v |

| Quantity: | 100 μg |
|----------------------|---|
| Target: | BHLHE41 |
| Binding Specificity: | AA 185-285 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BHLHE41 antibody is conjugated to Biotin |
| Application: | ELISA |

Product Details

| Immunogen: | Recombinant Human Class E basic helix-loop-helix protein 41 protein (185-285AA) |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | BHLHE41 |
|-------------------|---|
| Alternative Name: | BHLHE41 (BHLHE41 Products) |
| Background: | Background: Transcriptional repressor involved in the regulation of the circadian rhythm by |
| | negatively regulating the activity of the clock genes and clock-controlled genes. Acts as the |

negative limb of a novel autoregulatory feedback loop (DEC loop) which differs from the one formed by the PER and CRY transcriptional repressors (PER/CRY loop). Both these loops are interlocked as it represses the expression of PER1 and in turn is repressed by PER1/2 and CRY1/2. Represses the activity of the circadian transcriptional activator: CLOCK-ARNTL/BMAL1 heterodimer by competing for the binding to E-box elements (5\'-CACGTG-3\') found within the promoters of its target genes. Negatively regulates its own expression and the expression of DBP and BHLHE41/DEC2. Acts as a corepressor of RXR and the RXR-LXR heterodimers and represses the ligand-induced RXRA/B/G, NR1H3/LXRA, NR1H4 and VDR transactivation activity.

Aliases: Basic helix-loop-helix domain containing class B3 antibody, BHE41_HUMAN antibody, bHLH protein DEC2 antibody, bHLHB3 antibody, bHLHe41 antibody, Class B basic helix loop helix protein 3 antibody, Class B basic helix-loop-helix protein 3 antibody, Class E basic helixloop-helix protein 41 antibody, DEC2 antibody, Differentially expressed in chondrocytes protein 2 antibody, Enhancer of split and hairy related protein 1 antibody, Enhancer-of-split and hairyrelated protein 1 antibody, hDEC2 antibody, Moderately similar to basic-helix loop helix protein M.musculus antibody, SHARP-1 antibody

UniProt: Q9C0J9

Pathways: Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Buffer:

Preservative:

Format: Liquid

Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

handled by trained staff only.

Preservative: 0.03 % Proclin 300

-20 °C,-80 °C Storage:

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.