# antibodies .- online.com





# anti-COPB1 antibody (AA 2-200) (Biotin)



$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μg
Target:	COPB1
Binding Specificity:	AA 2-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This COPB1 antibody is conjugated to Biotin
Application:	ELISA

### **Product Details**

Immunogen:	Recombinant Human Coatomer subunit beta protein (2-200AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

# Target Details

Target:	COPB1
Alternative Name:	COPB1 (COPB1 Products)
Background:	Background: The coatomer is a cytosolic protein complex that binds to dilysine motifs and
	reversibly associates with Golgi non-clathrin-coated vesicles, which further mediate

biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins, the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors. Plays a functional role in facilitating the transport of kappa-type opioid receptor mRNAs into axons and enhances translation of these proteins. Required for limiting lipid storage in lipid droplets. Involved in lipid homeostasis by regulating the presence of perilipin family members PLIN2 and PLIN3 at the lipid droplet surface and promoting the association of adipocyte surface triglyceride lipase (PNPLA2) with the lipid droplet to mediate lipolysis (By similarity). Involved in the Golgi disassembly and reassembly processes during cell cycle. Involved in autophagy by playing a role in early endosome function. Plays a role in organellar compartmentalization of secretory compartments including endoplasmic reticulum (ER)-Golgi intermediate compartment (ERGIC), Golgi, trans-Golgi network (TGN) and recycling endosomes, and in biosynthetic transport of CAV1. Promotes degradation of Nef cellular targets CD4 and MHC class I antigens by facilitating their trafficking to degradative compartments.

Aliases: Beta-coat protein antibody, Beta-COP antibody, betacop antibody, Coatomer beta subunit antibody, Coatomer protein complex subunit beta 1 antibody, Coatomer protein complex subunit beta antibody, Coatomer subunit beta antibody, COPB antibody, COPB\_HUMAN antibody, Copb1 antibody, DKFZp761K102 antibody, FLJ10341 antibody

UniProt:

P53618

#### **Application Details**

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

# Handling

	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.