antibodies -online.com





anti-COPE antibody (AA 2-308) (HRP)



Go to Product page

| () | 11/0 | K\ / | iew |
|-----|--------|--------|-------------|
| | \cup | 'I V/I | $I \cap VV$ |
| | | | |

| Quantity: | 100 μg |
|----------------------|---|
| Target: | COPE |
| Binding Specificity: | AA 2-308 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This COPE antibody is conjugated to HRP |
| Application: | ELISA |

Product Details

| Immunogen: | Recombinant Human Coatomer subunit epsilon protein (2-308AA) | |
|-------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Purification: | >95%, Protein G purified | |

Target Details

| Target: | COPE |
|-------------------|---|
| Alternative Name: | COPE (COPE Products) |
| Background: | Background: The coatomer is a cytosolic protein complex that binds to dilysine motifs and |
| | reversibly associates with Golgi non-clathrin-coated vesicles, which further mediate |

Target Details

biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. The coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated with ADP-ribosylation factors (ARFs), which are small GTP-binding proteins, the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors.

Aliases: 1110005D17Rik antibody, Coatomer epsilon subunit antibody, Coatomer protein complex subunit epsilon antibody, Coatomer subunit epsilon antibody, COPE antibody, COPE_HUMAN antibody, Cope1 antibody, Epsilon coat protein antibody, Epsilon COP antibody, Epsilon COP I antibody, Epsilon subunit of coatomer protein complex antibody, Epsilon-coat protein antibody, Epsilon-COP antibody, FLJ13241 antibody

UniProt:

014579

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 |
| | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be |
| | handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |