# antibodies - online.com







## anti-CCDC6 antibody (AA 347-440)





-						
	\/		r١	/1	0	۱۸/
$\sim$	V	$\overline{}$	1 V	١,	$\sim$	VΥ

Quantity:	100 μg
Target:	CCDC6
Binding Specificity:	AA 347-440
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC6 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC)

#### **Product Details**

Immunogen:	Recombinant Human Coiled-coil domain-containing protein 6 protein (347-440AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	CCDC6	
Alternative Name:	CCDC6 (CCDC6 Products)	
Background:	Background: cytosol, structural constituent of cytoskeleton	
	Aliases: CCDC 6 antibody, CCDC6 antibody, CCDC6_HUMAN antibody, Coiled coil domain	

#### **Target Details**

containing 6 antibody, Coiled coil domain containing protein 6 antibody, Coiled-coil domain-containing protein 6 antibody, D10S170 antibody, FLJ32286 antibody, H4 antibody, Papillary thyroid carcinoma encoded protein antibody, Papillary thyroid carcinoma-encoded protein antibody, Protein H4 antibody, PTC antibody, TPC antibody, TST 1 antibody, TST1 antibody

UniProt:

Q16204


#### **Application Details**

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:500-1:1000, IF:1:50-1:200,
Restrictions:	For Research Use only

#### Handling


Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

#### **Images**




#### Immunohistochemistry

Image 1. IHC image of ABIN7148132 diluted at 1:500 and staining in paraffin-embedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



#### **Western Blotting**

**Image 2.** Western Blot Positive WB detected in: MCF-7 whole cell lysate All lanes: CCDC6 antibody at 3 μg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 54 kDa Observed band size: 54 kDa



#### **Immunohistochemistry**

Image 3. IHC image of ABIN7148132 diluted at 1:500 and staining in paraffin-embedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Please check the product details page for more images. Overall 4 images are available for ABIN7148132.