antibodies

# Datasheet for ABIN7148152 anti-CCDC81 antibody (AA 228-323)

I Image



## Overview

Quantity:	100 µg
Target:	CCDC81
Binding Specificity:	AA 228-323
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC81 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

## Product Details

Immunogen:	Recombinant Human Coiled-coil domain-containing protein 81 protein (228-323AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details


Target:	CCDC81
Alternative Name:	CCDC81 (CCDC81 Products)
Background:	Background: centrosome
	Aliases: CCD81_HUMAN antibody, Ccdc81 antibody, Coiled-coil domain containing 81 antibody,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7148152 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

## Target Details

	Coiled-coil domain-containing protein 81 antibody	
UniProt:	Q6ZN84	
Application Details		
Application Notes:	Recommended dilution: IHC:1:200-1:500,	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300	
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	

## Images



#### Immunohistochemistry

**Image 1.** IHC image of ABIN7148152 diluted at 1:400 and staining in paraffin-embedded human pancreatic cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.