



[Go to Product page](#)

Datasheet for ABIN7148184
anti-CHCHD5 antibody (AA 1-110)

2 Images

Overview

Quantity:	100 µL
Target:	CHCHD5
Binding Specificity:	AA 1-110
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CHCHD5 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA

Product Details

Immunogen:	Recombinant Human Coiled-coil-helix-coiled-coil-helix domain-containing protein 5 protein (1-110AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CHCHD5
Alternative Name:	CHCHD5 (CHCHD5 Products)
Background:	Background: mitochondrial intermembrane space, aerobic respiration

Target Details

Aliases: 1110027L01Rik antibody, AW045710 antibody, C2orf9 antibody, CHCH5_HUMAN antibody, CHCHD5 antibody, chromosome 2 open reading frame antibody, Coiled coil helix coiled coil helix domain containing 5 antibody, Coiled-coil-helix-coiled-coil-helix domain-containing protein 5 antibody, FLJ39671 antibody, MGC11104 antibody, RP24-388N11.2 antibody

UniProt: [Q9BSY4](#)

Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7148184 diluted at 1:200 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP



and visualized using 0.05 % DAB.

Immunohistochemistry

Image 2. IHC image of ABIN7148184 diluted at 1:200 and staining in paraffin-embedded human ovarian cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05 % DAB.