

Datasheet for ABIN7148345

anti-C1QTNF3 antibody (AA 23-246)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	C1QTNF3
Binding Specificity:	AA 23-246
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C1QTNF3 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Complement C1q tumor necrosis factor-related protein 3 protein (23-246AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	C1QTNF3
Alternative Name:	C1QTNF3 (C1QTNF3 Products)
Background:	Background: C1QTNF3 is a secretory protein produced by chondrogenic precursors &

Target Details

proliferating chondrocytes, and belongs to a novel C1q family of proteins.

Aliases: C1ATNF3 antibody, C1q and tumor necrosis factor related protein 3 antibody, C1QT3_HUMAN antibody, C1qtnf3 antibody, Cartonectin antibody, Collagenous repeat containing sequence of 26 kDa antibody, collagenous repeat-containing sequence 26 kDa protein antibody, Complement C1q tumor necrosis factor related protein 3 antibody, Complement C1q tumor necrosis factor-related protein 3 antibody, Corcs antibody, CORS 26 antibody, Cors antibody, CORS26 antibody, CTRP3 antibody, FLJ37576 antibody, OTTHUMP00000115918 antibody, OTTHUMP00000219981 antibody, PRO1484 antibody, Secretory protein CORS26 antibody, UNQ753 antibody

UniProt: [Q9BXJ4](#)

Application Details

Application Notes: Recommended dilution: IHC:1:20-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7148345 diluted at 1:100 and staining in paraffin-embedded human small intestine tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.