

Datasheet for ABIN7148466  
**anti-COG8 antibody (AA 521-612)**[Go to Product page](#)

## 1 Image

## Overview

|                      |                                     |
|----------------------|-------------------------------------|
| Quantity:            | 100 µg                              |
| Target:              | COG8                                |
| Binding Specificity: | AA 521-612                          |
| Reactivity:          | Human                               |
| Host:                | Rabbit                              |
| Clonality:           | Polyclonal                          |
| Conjugate:           | This COG8 antibody is un-conjugated |
| Application:         | ELISA, Immunohistochemistry (IHC)   |

## Product Details

|                   |  |
|-------------------|--|
| Immunogen:        | Recombinant Human Conserved oligomeric Golgi complex subunit 8 protein (521-612AA) |
| Isotype:          | IgG  |
| Cross-Reactivity: | Human  |
| Purification:     | >95%, Protein G purified   |

## Target Details

|                   |   |
|-------------------|---|
| Target:           | COG8  |
| Alternative Name: | COG8 ( <a href="#">COG8 Products</a> )  |
| Background:       | Background: Required for normal Golgi function.<br>Aliases: COG8 antibody, Conserved oligomeric Golgi complex subunit 8 antibody, COG complex |

## Target Details

subunit 8 antibody, Component of oligomeric Golgi complex 8 antibody

UniProt: [Q96MW5](#)

## Application Details

Application Notes: Recommended dilution: IHC:1:20-1:200,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



### Immunohistochemistry

**Image 1.** IHC image of ABIN7148466 diluted at 1:100 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.