-online.com antibodies

## Datasheet for ABIN7148563 anti-H2AFY2 antibody (AA 1-260) (Biotin)



Overview

| Quantity:            | 100 µg                                       |
|----------------------|--|
| Target:              | H2AFY2                                       |
| Binding Specificity: | AA 1-260                                     |
| Reactivity:          | Human  |
| Host:                | Rabbit                                       |
| Clonality:           | Polyclonal                                   |
| Conjugate:           | This H2AFY2 antibody is conjugated to Biotin |
| Application:         | ELISA  |

## Product Details

| Immunogen:        | Recombinant Human Core histone macro-H2A.2 protein (1-260AA) |
|-------------------|--|
| Isotype:          | lgG  |
| Cross-Reactivity: | Human  |
| Purification:     | >95%, Protein G purified                                     |

## Target Details

| Target:           | H2AFY2  |
|-------------------|---|
| Alternative Name: | H2AFY2 (H2AFY2 Products)  |
| Background:       | Background: Variant histone H2A which replaces conventional H2A in a subset of nucleosomes  |
|                   | where it represses transcription. Nucleosomes wrap and compact DNA into chromatin, limiting |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7148563 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

|                     | DNA accessibility to the cellular machineries which require DNA as a template. Histones       |
|---------------------|---|
|                     | thereby play a central role in transcription regulation, DNA repair, DNA replication and      |
|                     | chromosomal stability. DNA accessibility is regulated via a complex set of post-translational |
|                     | modifications of histones, also called histone code, and nucleosome remodeling. May be        |
|                     | involved in stable X chromosome inactivation.   |
|                     | Aliases: Core histone macro H2A2.2 antibody, Core histone macro-H2A.2 antibody, Core          |
|                     | histone macroH2A2.2 antibody, H2A histone family member Y2 antibody, H2AFY2 antibody,         |
|                     | H2AW_HUMAN antibody, Histone macroH2A2 antibody, Macro H2A.2 antibody, Macro H2A2             |
|                     | antibody, MacroH2A.2 antibody, MacroH2A2 antibody, mH2A2 antibody                             |
| UniProt:            | Q9P0M6  |
| Application Details |   |
| Application Notes:  | Optimal working dilution should be determined by the investigator.                            |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Buffer:             | Preservative: 0.03 % Proclin 300  |
|                     | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should b             |
|                     | handled by trained staff only.  |
| Storage:            | -20 °C,-80 °C   |
| Storage Comment:    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                                 |
|                     |   |