antibodies.com

Datasheet for ABIN7149159 anti-COX2 antibody (AA 83-227) (Biotin)



\sim			
()	ver	1/1C	۱۸/
\cup	マロ	VIC	7 V V

Quantity:	100 µg
Target:	COX2
Binding Specificity:	AA 83-227
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This COX2 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Cytochrome c oxidase subunit 2 protein (83-227AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	COX2
Alternative Name:	MT-CO2 (COX2 Products)
Background: Background: Cytochrome c oxidase is the component of the respiratory chain that c	
	reduction of oxygen to water. Subunits 1-3 form the functional core of the enzyme complex.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7149159 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Subunit 2 transfers the electrons from cytochrome c via its binu bimetallic center of the catalytic subunit 1. Aliases: COII antibody, COX 2 antibody, COX II antibody, COX2 ar antibody, COXII antibody, COX 2 antibody, COX II antibody, COX2 ar antibody, COXII antibody, Cytochrome c oxidase II antibody, Cyto II antibody, Cytochrome c oxidase subunit 2 antibody, MT CO2 a MTCO2 antibody UniProt: P00403 Pathways: Brown Fat Cell Differentiation, Positive Regulation of fat Cell Diff Application Details Application Notes: Optimal working dilution should be determined by the investigat Restrictions: For Research Use only Handling	ntibody, COX2_HUMAN ochrome c oxidase polypeptide ntibody, MT-CO2 antibody, erentiation	
Aliases: COII antibody, COX 2 antibody, COX II antibody, COX2 ar antibody, COXII antibody, Cytochrome c oxidase II antibody, Cyto II antibody, Cytochrome c oxidase subunit 2 antibody, MT CO2 a MTCO2 antibodyUniProt:P00403Pathways:Brown Fat Cell Differentiation, Positive Regulation of fat Cell Diff Application DetailsApplication Notes:Optimal working dilution should be determined by the investigat For Research Use only	ochrome c oxidase polypeptide ntibody, MT-CO2 antibody, erentiation	
antibody, COXII antibody, Cytochrome c oxidase II antibody, Cyto II antibody, Cytochrome c oxidase subunit 2 antibody, MT CO2 a MTCO2 antibody UniProt: P00403 Pathways: Brown Fat Cell Differentiation, Positive Regulation of fat Cell Diff Application Details Optimal working dilution should be determined by the investigat Restrictions: For Research Use only	ochrome c oxidase polypeptide ntibody, MT-CO2 antibody, erentiation	
II antibody, Cytochrome c oxidase subunit 2 antibody, MT CO2 a MTCO2 antibody UniProt: P00403 Pathways: Brown Fat Cell Differentiation, Positive Regulation of fat Cell Diff Application Details Optimal working dilution should be determined by the investigat Restrictions: For Research Use only	ntibody, MT-CO2 antibody, erentiation	
MTCO2 antibody UniProt: P00403 Pathways: Brown Fat Cell Differentiation, Positive Regulation of fat Cell Diff Application Details Optimal working dilution should be determined by the investigat Restrictions: For Research Use only	erentiation	
UniProt:P00403Pathways:Brown Fat Cell Differentiation, Positive Regulation of fat Cell DiffApplication DetailsApplication Notes:Optimal working dilution should be determined by the investigatRestrictions:For Research Use only		
Pathways: Brown Fat Cell Differentiation, Positive Regulation of fat Cell Diff Application Details Application Notes: Optimal working dilution should be determined by the investigat Restrictions: For Research Use only		
Application Details Application Notes: Optimal working dilution should be determined by the investigat Restrictions: For Research Use only		
Application Notes: Optimal working dilution should be determined by the investigat Restrictions: For Research Use only	or.	
Restrictions: For Research Use only	Dr.	
Handling		
Format: Liquid		
Buffer: Preservative: 0.03 % Proclin 300	Preservative: 0.03 % Proclin 300	
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4		
Preservative: ProClin		
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS	SUBSTANCE which should be	
handled by trained staff only.		
Storage: -20 °C,-80 °C		
Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.		