



[Go to Product page](#)

Datasheet for ABIN7149210  
**anti-CYP1B1 antibody (AA 282-409)**

1 Image

### Overview

Quantity:	100 µg
Target:	CYP1B1
Binding Specificity:	AA 282-409
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP1B1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

### Product Details

Immunogen:	Recombinant Human Cytochrome P450 1B1 protein (282-409AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	CYP1B1
Alternative Name:	CYP1B1 ( <a href="#">CYP1B1 Products</a> )
Background:	Background: Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It

## Target Details

---

oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, retinoid and xenobiotics. Preferentially oxidizes 17beta-estradiol to the carcinogenic 4-hydroxy derivative, and a variety of procarcinogenic compounds to their activated forms, including polycyclic aromatic hydrocarbons. Promotes angiogenesis by removing cellular oxygenation products, thereby decreasing oxidative stress, release of antiangiogenic factor THBS2, then allowing endothelial cells migration, cell adhesion and capillary morphogenesis. These changes are concomitant with the endothelial nitric oxide synthase activity and nitric oxide synthesis. Plays an important role in the regulation of perivascular cell proliferation, migration, and survival through modulation of the intracellular oxidative state and NF-kappa-B expression and/or activity, during angiogenesis. Contributes to oxidative homeostasis and ultrastructural organization and function of trabecular meshwork tissue through modulation of POSTN expression.

Aliases: Aryl hydrocarbon hydroxylase antibody, CP1B antibody, CP1B1\_HUMAN antibody, Cyp1b1 antibody, CYPIB1 antibody, Cytochrome P450 1B1 antibody, Cytochrome P450 family 1 subfamily B polypeptide 1 antibody, Cytochrome P450 subfamily I (dioxin inducible) polypeptide 1 (glaucoma 3 primary infantile) antibody, Flavoprotein linked monooxygenase antibody, GLC3A antibody, Microsomal monooxygenase antibody, P4501B1 antibody, Xenobiotic monooxygenase antibody

---

UniProt: [Q16678](#)

---

Pathways: [Steroid Hormone Biosynthesis](#)

## Application Details

---

Application Notes: Recommended dilution: WB:1:500-1:5000,

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


---

## Handling

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



### Western Blotting

**Image 1.** Western Blot Positive WB detected in: PC-3 whole cell lysate, A549 whole cell lysate All lanes: CYP1B1 antibody at 6.62 µg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 61 kDa Observed band size: 61 kDa