# antibodies .- online.com







# anti-FAM96A antibody (AA 1-160) (Biotin)



| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:            | 100 μg                                       |
|----------------------|--|
| Target:              | FAM96A                                       |
| Binding Specificity: | AA 1-160                                     |
| Reactivity:          | Human  |
| Host:                | Rabbit                                       |
| Clonality:           | Polyclonal                                   |
| Conjugate:           | This FAM96A antibody is conjugated to Biotin |
| Application:         | ELISA  |

### **Product Details**

| Immunogen:        | Recombinant Human Cytosolic iron-sulfur assembly component 2A protein (1-160AA) |  |
|-------------------|---|--|
| Isotype:          | IgG   |  |
| Cross-Reactivity: | Human   |  |
| Purification:     | >95%, Protein G purified  |  |

# Target Details

| Target:           | FAM96A   |
|-------------------|--|
| Alternative Name: | FAM96A (FAM96A Products)   |
| Background:       | Background: Component of the cytosolic iron-sulfur protein assembly (CIA) complex, a |
|                   | multiprotein complex that mediates the incorporation of iron-sulfur cluster into     |

#### **Target Details**

extramitochondrial Fe/S proteins (PubMed:23891004). As a CIA complex component and in collaboration with CIAO1 specifically matures ACO1 and stabilizes IREB2, connecting cytosolic iron-sulfur protein maturation with cellular iron regulation (PubMed:23891004). May play a role in chromosome segregation through establishment of sister chromatid cohesion. May induce apoptosis in collaboration with APAF1 (PubMed:25716227)

Aliases: CIAO2A antibody, CIA2A antibody, FAM96A antibody, Cytosolic iron-sulfur assembly component 2A antibody, MIP18 family protein FAM96A antibody

UniProt:

Storage:

Storage Comment:

Q9H5X1

## **Application Details**

| Application Notes: | Optimal working dilution should be determined by the investigator.                 |  |
|--------------------|--|--|
| Restrictions:      | For Research Use only  |  |
|                    |  |  |
| Handling           |  |  |
| Format:            | Liquid   |  |
| Buffer:            | Preservative: 0.03 % Proclin 300   |  |
|                    | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4                                     |  |
| Preservative:      | ProClin  |  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be |  |

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

handled by trained staff only.

-20 °C,-80 °C