



[Go to Product page](#)

Datasheet for ABIN7149490
anti-DACT1 antibody (AA 410-525) (FITC)

Overview

Quantity:	100 µg
Target:	DACT1
Binding Specificity:	AA 410-525
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DACT1 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Dapper homolog 1 protein (410-525AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	DACT1
Alternative Name:	DACT1 (DACT1 Products)
Background:	Background: Involved in regulation of intracellular signaling pathways during development. Specifically thought to play a role in canonical and/or non-canonical Wnt signaling pathways

Target Details

through interaction with DSH (Dishevelled) family proteins. The activation/inhibition of Wnt signaling may depend on the phosphorylation status. Proposed to regulate the degradation of CTNNB1/beta-catenin, thereby modulating the transcriptional activation of target genes of the Wnt signaling pathway. Its function in stabilizing CTNNB1 may involve inhibition of GSK3B activity. Promotes the membrane localization of CTNNB1. The cytoplasmic form can induce DVL2 degradation via a lysosome-dependent mechanism, the function is inhibited by PKA-induced binding to 14-3-3 proteins, such as YWHAB. Seems to be involved in morphogenesis at the primitive streak by regulating VANGL2 and DVL2, the function seems to be independent of canonical Wnt signaling and rather involves the non-canonical Wnt/planar cell polarity (PCP) pathway (By similarity). The nuclear form may prevent the formation of LEF1:CTNNB1 complex and recruit HDAC1 to LEF1 at target gene promoters to repress transcription thus antagonizing Wnt signaling. May be involved in positive regulation of fat cell differentiation. During neuronal differentiation may be involved in excitatory synapse organization, and dendrite formation and establishment of spines.

Aliases: Dact antibody, dact1 antibody, DACT1_HUMAN antibody, Dapper antagonist of beta catenin homolog 1 antibody, Dapper antagonist of catenin 1 antibody, Dapper antibody, Dapper homolog 1 antibody, DAPPER1 antibody, Dpr antibody, Dpr1 antibody, Frd antibody, Frodo antibody, hDPR1 antibody, Hepatocellular carcinoma novel gene 3 protein antibody, Heptacellular carcinoma novel gene 3 antibody, HNG3 antibody, MTNG3 antibody, Thyex3 antibody

UniProt: [Q9NYF0](#)

Pathways: [Positive Regulation of fat Cell Differentiation](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.