

Datasheet for ABIN7149571
anti-PDSS2 antibody (AA 292-387)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	PDSS2
Binding Specificity:	AA 292-387
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDSS2 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Decaprenyl-diphosphate synthase subunit 2 protein (292-387AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	PDSS2
Alternative Name:	PDSS2 (PDSS2 Products)
Background:	Background: Supplies decaprenyl diphosphate, the precursor for the side chain of the isoprenoid quinones ubiquinone-10.

Target Details

Aliases: All-trans-decaprenyl-diphosphate synthase subunit 2 antibody, bA59I9.3 antibody, C6orf210 antibody, Candidate tumor suppressor protein antibody, chromosome 6 open reading frame 210 antibody, Decaprenyl pyrophosphate synthase subunit 2 antibody, decaprenyl pyrophosphate synthetase subunit 2 antibody, Decaprenyl-diphosphate synthase subunit 2 antibody, DLP1 antibody, DLP1_HUMAN antibody, hDLP1 antibody, Pdss2 antibody, prenyl (decaprenyl) diphosphate synthase, subunit 2 antibody, subunit 2 of decaprenyl diphosphate synthase antibody

UniProt: [Q86YH6](#)

Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7149571 diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7149571 diluted at 1:200 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.