antibodies .- online.com





anti-DOCK7 antibody (AA 1440-1671) (Biotin)



\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μg
Target:	DOCK7
Binding Specificity:	AA 1440-1671
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DOCK7 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Dedicator of cytokinesis protein 7 protein (1440-1671AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	DOCK7
Alternative Name:	DOCK7 (DOCK7 Products)
Background:	Background: Functions as a guanine nucleotide exchange factor (GEF), which activates Rac1
	and Rac3 Rho small GTPases by exchanging bound GDP for free GTP. Does not have a GEF

Target Details

activity for CDC42. Required for STMN1 \\\'Ser-15\\\' phosphorylation during axon formation and consequently for neuronal polarization (PubMed:16982419). Has a role in pigmentation (By similarity). Involved in the regulation of cortical neurogenesis through the control of radial glial cells (RGCs) proliferation versus differentiation, negatively regulates the basal-to-apical interkinetic nuclear migration of RGCs by antagonizing the microtubule growth-promoting function of TACC3 (By similarity).

Aliases: DOCK7 antibody, KIAA1771Dedicator of cytokinesis protein 7 antibody

UniProt:

Storage:

Storage Comment:

Q96N67

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

handled by trained staff only.

-20 °C,-80 °C