# antibodies - online.com







# anti-DHRS9 antibody (AA 23-319)

**Images** 



#### Overview

Quantity:	100 μg
Target:	DHRS9
Binding Specificity:	AA 23-319
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DHRS9 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

#### **Product Details**

Immunogen:	Recombinant Human Dehydrogenase/reductase SDR family member 9 protein (23-319AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	DHRS9
Alternative Name:	DHRS9 (DHRS9 Products)
Background:	Background: Involved in the targeting and/or fusion of transport vesicles to their target
	membrane during transport of proteins from the early endosome to the lysosome. Required for

heterotypic fusion of late endosomes with lysosomes and homotypic lysosomal fusion.

Required for calcium regulated lysosomal exocytosis. Involved in the export of chylomicrons from the endoplasmic reticulum to the cis Golgi. Required for exocytosis of mediators during eosinophil and neutrophil degranulation, and target cell killing by natural killer cells. Required for focal exocytosis of late endocytic vesicles during phagosome formation.

Aliases: 3 alpha hydroxysteroid dehydrogenase antibody, 3-@alpha-HSD antibody, 3-@alpha-hydroxysteroid dehydrogenase antibody, 3-alpha hydroxysteroid dehydrogenase antibody, 3-alpha-HSD antibody, 3alpha HSD antibody, Dehydrogenase/reductase SDR family member 9 antibody, DHRS9 antibody, DHRS9\_HUMAN antibody, NADP dependent retinol dehydrogenase/reductase antibody, NADP-dependent retinol dehydrogenase/reductase antibody, RDH-E2 antibody, RDHL antibody, RDHTBE antibody, retinol dehydrogenase homolog antibody, Retinol dehydrogenase, tracheobronchiol epithelail cell-specific antibody, RETSDR8 antibody, SDR family, member 9 antibody, SDR9C4 antibody, short chain dehydrogenase/reductase family 9C, member 4 antibody, Short chain dehydrogenase/reductase retSDR8 antibody, Short-chain dehydrogenase/reductase retSDR8 antibody

UniProt: Q9BPW9

Pathways: C21-Steroid Hormone Metabolic Process

#### **Application Details**

Application Notes: Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200,

Restrictions: For Research Use only


#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C

Storage Comment:


Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

### **Images**



#### **Immunohistochemistry**

**Image 1.** Immunohistochemistry of paraffin-embedded human kidney tissue using ABIN7149627 at dilution of 1:100



#### **Immunofluorescence**

**Image 2.** Immunofluorescent analysis of A549 cells using ABIN7149627 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)