antibodies - online.com







anti-DHRSX antibody (AA 32-330)

Images



| \sim | | |
|--------|-------|--------------------|
| () | IV/Ar | $V_{i} \cap V_{i}$ |

| Overview | |
|----------------------|------------------------------------------------------------------------------------------------|
| Quantity: | 100 μg |
| Target: | DHRSX |
| Binding Specificity: | AA 32-330 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This DHRSX antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF) |
| Product Details | |
| Immunogen: | Recombinant Human Dehydrogenase/reductase SDR family member on chromosome X protein (32-330AA) |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | DHRSX |
|-------------------|---------------------------------------------------------------------------|
| Alternative Name: | DHRSX (DHRSX Products) |
| Background: | CXorf11 antibody, Dehydrogenase/reductase (SDR family) X linked antibody, |

Target Details

| Dehydrogenase/reductase SDR family member on chromosome X antibody, DHRS5X antibo | ody, |
|-----------------------------------------------------------------------------------|------|
| DHRS5Y antibody, DHRSX antibody, DHRSX_HUMAN antibody, DHRSXY antibody, DHRSY | |
| antibody, RP11 325D5.2 antibody | |

UniProt:

Q8N5I4


Application Details

| Application Notes: | Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200, |
|--------------------|------------------------------------------------------|
| Restrictions: | For Research Use only |

Handling


| Format: | Liquid | |
|--------------------|-------------------------------------------------------------------------------------------------------------------|--|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | -20 °C,-80 °C | |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. | |

Images



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human adrenal gland tissue using ABIN7149631 at dilution of 1:100



Immunofluorescence

Image 2. Immunofluorescence staining of A549 cells with ABIN7149631 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).