antibodies -online.com





anti-Transmembrane 7 Superfamily Member 4 (TM7SF4) (AA 119-209) antibody



Go to Product pag

3 Images

Overview	
Quantity:	100 μg
Target:	Transmembrane 7 Superfamily Member 4 (TM7SF4)
Binding Specificity:	AA 119-209
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Un-conjugated
Application:	ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC)
Product Details	
Immunogen:	Recombinant Human Dendritic cell-specific transmembrane protein (119-209AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified
Target Details	
Target:	Transmembrane 7 Superfamily Member 4 (TM7SF4)
Alternative Name:	DCSTAMP (TM7SF4 Products)
Background:	Background: Probable cell surface receptor that plays several roles in cellular fusion, cell

differentiation, bone and immune homeostasis. Plays a role in TNFSF11-mediated

osteoclastogenesis. Cooperates with OCSTAMP in modulating cell-cell fusion in both osteoclasts and foreign body giant cells (FBGCs). Participates in osteoclast bone resorption. Involved in inducing the expression of tartrate-resistant acid phosphatase in osteoclast precursors. Plays a role in haematopoietic stem cell differentiation of bone marrow cells toward the myeloid lineage. Inhibits the development of neutrophilic granulocytes. Plays also a role in the regulation of dendritic cell (DC) antigen presentation activity by controlling phagocytic activity. Involved in the maintenance of immune self-tolerance and avoidance of autoimmune reactions.

Aliases: DC-STAMP antibody, DCSTAMP antibody, DCSTP_HUMAN antibody, Dendritic cell-specific transmembrane protein antibody, Dendrocyte-expressed seven transmembrane protein antibody, FIND antibody, hDC-STAMP antibody, IL-Four INDuced antibody, IL-four-induced protein antibody, Transmembrane 7 superfamily member 4 antibody

UniProt:

Q9H295

Application Details

Storage Comment:

Application Notes:	Recommended dilution: IHC:1:200-1:500, IF:1:50-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid


Buffer:
Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative:
ProClin

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunohistochemistry

Image 1. IHC image of ABIN7149663 diluted at 1:200 and staining in paraffin-embedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence

Image 2. Immunofluorescence staining of HepG2 cells with ABIN7149663 at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemistry

Image 3. IHC image of ABIN7149663 diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.