antibodies - online.com







anti-DDA1 antibody (AA 1-102)





\sim		
()\/ (rview	1
\circ	IVICV	٧

Quantity:	100 μg
Target:	DDA1
Binding Specificity:	AA 1-102
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DDA1 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human DET1- and DDB1-associated protein 1 protein (1-102AA)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	>95%, Protein G purified

Target Details

Target:	DDA1
Alternative Name:	DDA1 (DDA1 Products)
Background:	Background: May be involved in ubiquitination and subsequent proteasomal degradation of
	target proteins. Component of the DDD-E2 complexes which may provide a platform for

interaction with CUL4A and WD repeat proteins.

Aliases: C19orf58 antibody, Cross immune reaction antigen PCIA1 antibody, DDA1 antibody, DDA1_HUMAN antibody, DET1 and DDB1 associated 1 antibody, DET1 and DDB1 associated protein 1 antibody, DET1- and DDB1-associated protein 1 antibody, MGC2594 antibody, PCIA-1 antibody, PCIA1 antibody, Placenta cross immune reaction antigen 1 antibody, Placenta cross-immune reaction antigen 1 antibody

UniProt:

Q9BW61

Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid


Buffer:
Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative:
ProClin

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunofluorescence

Image 1. Immunofluorescent analysis of PC-3 cells using ABIN7149804 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western Blotting

Image 2. Western Blot Positive WB detected in: Rat liver tissue, Mouse liver tissue All lanes: DDA1 antibody at 2.8 μ g/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 12 kDa Observed band size: 12 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human kidney tissue using ABIN7149804 at dilution of 1:100