

## Datasheet for ABIN7149821 anti-DPPA3 antibody (AA 1-159) (Biotin)



Overview

Quantity:	100 µg
Target:	DPPA3
Binding Specificity:	AA 1-159
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DPPA3 antibody is conjugated to Biotin
Application:	ELISA

## Product Details

Immunogen:	Recombinant Human Developmental pluripotency-associated protein 3 protein (1-159AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	DPPA3
Alternative Name:	DPPA3 (DPPA3 Products)
Background:	Background: Primordial germ cell (PGCs)-specific protein involved in epigenetic chromatin reprogramming in the zygote following fertilization. In zygotes, DNA demethylation occurs

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7149821 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

antibody, Stella related protein antibody, Stella-related protein antibody
pluripotency-associated protein 3 antibody, Dppa3 antibody, DPPA3_HUMAN antibody, PGC7
antibody, Developmental pluripotency associated protein 3 antibody, Developmental
Aliases: Compaction associated protein 1 antibody, Developmental pluripotency associated 3
development. Also involved in chromatin condensation in oocytogenesis (By similarity).
May be important for the totipotent/pluripotent states continuing through preimplantation
are marked with H3K9me2 in mature sperm from DNA demethylation in early embryogenesis.
into protamine and does not contain much H3K9me2 mark. Also protects imprinted loci that
and subsequent DNA demethylation. Does not bind paternal chromatin, which is mainly packed
maternal genome, and protects maternal genome from TET3-mediated conversion to 5hmC
5hmC: specifically recognizes and binds histone H3 dimethylated at \'Lys-9\' (H3K9me2) on
protection of DNA methylation in the maternal pronucleus by preventing conversion of 5mC to
pronucleus and certain paternally-imprinted loci are protected from this process. Participates in
selectively in the paternal pronucleus before the first cell division, while the adjacent maternal

UniProt:

Q6W0C5

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.