

## Datasheet for ABIN7149987

# anti-DPP4 antibody (AA 473-588) (HRP)



_					
	W	0	rv	10	W

Quantity:	100 μg
Target:	DPP4
Binding Specificity:	AA 473-588
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DPP4 antibody is conjugated to HRP
Application:	ELISA

### **Product Details**

Immunogen:	Recombinant Human Dipeptidyl peptidase 4 protein (473-588AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### **Target Details**

Target:	DPP4
Alternative Name:	DPP4 (DPP4 Products)
Background:	Background: Cell surface glycoprotein receptor involved in the costimulatory signal essential for
	T-cell receptor (TCR)-mediated T-cell activation. Acts as a positive regulator of T-cell

coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC. Its binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3dependent manner. Its interaction with ADA also regulates lymphocyte-epithelial cell adhesion. In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM. May be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation. When overexpressed, enhanced cell proliferation, a process inhibited by GPC3. Acts also as a serine exopeptidase with a dipeptidyl peptidase activity that regulates various physiological processes by cleaving peptides in the circulation, including many chemokines, mitogenic growth factors, neuropeptides and peptide hormones. Removes N-terminal dipeptides sequentially from polypeptides having unsubstituted N-termini provided that the penultimate residue is proline. Aliases: ADABP antibody, ADCP-2 antibody, ADCP2 antibody, Adenosine deaminase complexing protein 2 antibody, CD26 antibody, Dipeptidyl peptidase 4 antibody, Dipeptidyl peptidase 4 soluble form antibody, Dipeptidyl peptidase IV antibody, Dipeptidyl peptidase IV membrane form antibody, Dipeptidyl peptidase IV soluble form antibody, DPP IV antibody, DPP4 antibody, DPP4\_HUMAN antibody, DPPIV antibody, T-cell activation antigen CD26 antibody, TP103 antibody

UniProt: P27487

Pathways: Peptide Hormone Metabolism, Regulation of Leukocyte Mediated Immunity

#### **Application Details**

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C

Storage Comment:

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.