

Datasheet for ABIN7149989
anti-DPP9 antibody (AA 289-437)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	DPP9
Binding Specificity:	AA 289-437
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DPP9 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA

Product Details

Immunogen:	Recombinant Human Dipeptidyl peptidase 9 protein (289-437AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	DPP9
Alternative Name:	DPP9 (DPP9 Products)
Background:	Background: Dipeptidyl peptidase that cleaves off N-terminal dipeptides from proteins having a Pro or Ala residue at position 2.

Target Details

Aliases: Dipeptidyl peptidase 9 antibody, Dipeptidyl peptidase IV related protein 2 antibody, Dipeptidyl peptidase IV-related protein 2 antibody, Dipeptidyl peptidase IX antibody, Dipeptidyl peptidase like protein 9 antibody, Dipeptidyl peptidase-like protein 9 antibody, Dipeptidylpeptidase 9 antibody, Dipeptidylpeptidase IX antibody, DKFZp762F117 antibody, DP 9 antibody, DP9 antibody, DPLP 9 antibody, DPLP9 antibody, DPP 9 antibody, DPP IX antibody, DPP9 antibody, DPP9_HUMAN antibody, DPRP 2 antibody, DPRP-2 antibody, DPRP2 antibody, FLJ16073 antibody

UniProt: [Q86T12](#)

Application Details

Application Notes: Recommended dilution: IHC:1:500-1:1000,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7149989 diluted at 1:800 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.