Datasheet for ABIN7150082 anti-DLG5 antibody (AA 1580-1704) (FITC)

antibodies .-online.com



Overview

| Quantity: | 100 µL |
|----------------------|--|
| Target: | DLG5 |
| Binding Specificity: | AA 1580-1704 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This DLG5 antibody is conjugated to FITC |
| Application: | Please inquire |

Product Details

| Immunogen: | Recombinant Human Disks large homolog 5 protein (1580-1704AA) |
|-------------------|---|
| Isotype: | lgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | DLG5 |
|-------------------|---|
| Alternative Name: | DLG5 (DLG5 Products) |
| Background: | Background: Acts as a regulator of the Hippo signaling pathway (PubMed:28087714, |
| | PubMed:28169360). Negatively regulates the Hippo signaling pathway by mediating the |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7150082 | 09/10/2023 | Copyright antibodies-online. All rights reserved. interaction of MARK3 with STK3/4, bringing them together to promote MARK3-dependent hyperphosphorylation and inactivation of STK3 kinase activity toward LATS1 (PubMed:28087714). Positively regulates the Hippo signaling pathway by mediating the interaction of SCRIB with STK4/MST1 and LATS1 which is important for the activation of the Hippo signaling pathway. Involved in regulating cell proliferation, maintenance of epithelial polarity, epithelial-mesenchymal transition (EMT), cell migration and invasion (PubMed:28169360). Plays an important role in dendritic spine formation and synaptogenesis in cortical neurons, regulates synaptogenesis by enhancing the cell surface localization of Ncadherin. Acts as a positive regulator of hedgehog (Hh) signaling pathway. Plays a critical role in the early point of the SMO activity cycle by interacting with SMO at the ciliary base to induce the accumulation of KIF7 and GLI2 at the ciliary tip for GLI2 activation (By similarity). Aliases: Discs large homolog 5 (Drosophila) antibody, Discs large homolog 5 antibody, Discs large protein LP DLG antibody, Discs large protein P dlg antibody, Discs large protein P-dlg. antibody, Disks large homolog 5 antibody, DLG 5 antibody, DLG5 antibody, DLG5_HUMAN antibody, KIAA0583 antibody, Large type of P DLG antibody, LP DLG antibody, P DLG5 antibody, PDLG antibody, Placenta and prostate DLG antibody

UniProt:

Q8TDM6

Application Details

| Restrictions: | For Research Use only |
|--------------------|---|
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN7150082 | 09/10/2023 | Copyright antibodies-online. All rights reserved.