antibodies -online.com





anti-DLG5 antibody (AA 1580-1704) (HRP)



\sim	
()\/△	$r_{1}/1 \triangle_{1} \lambda_{1}$
\cup \vee \subset	rview

Quantity:	100 μL
Target:	DLG5
Binding Specificity:	AA 1580-1704
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DLG5 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Disks large homolog 5 protein (1580-1704AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	DLG5
Alternative Name:	DLG5 (DLG5 Products)
Background:	Background: Acts as a regulator of the Hippo signaling pathway (PubMed:28087714,
	PubMed:28169360). Negatively regulates the Hippo signaling pathway by mediating the

interaction of MARK3 with STK3/4, bringing them together to promote MARK3-dependent hyperphosphorylation and inactivation of STK3 kinase activity toward LATS1 (PubMed:28087714). Positively regulates the Hippo signaling pathway by mediating the interaction of SCRIB with STK4/MST1 and LATS1 which is important for the activation of the Hippo signaling pathway. Involved in regulating cell proliferation, maintenance of epithelial polarity, epithelial-mesenchymal transition (EMT), cell migration and invasion (PubMed:28169360). Plays an important role in dendritic spine formation and synaptogenesis in cortical neurons, regulates synaptogenesis by enhancing the cell surface localization of Ncadherin. Acts as a positive regulator of hedgehog (Hh) signaling pathway. Plays a critical role in the early point of the SMO activity cycle by interacting with SMO at the ciliary base to induce the accumulation of KIF7 and GLI2 at the ciliary tip for GLI2 activation (By similarity). Aliases: Discs large homolog 5 (Drosophila) antibody, Discs large homolog 5 antibody, Discs large protein LP DLG antibody, Discs large protein P dlg antibody, Discs large protein P-dlg antibody, Disks large homolog 5 antibody, DLG 5 antibody, DLG5 antibody, DLG5_HUMAN antibody, KIAA0583 antibody, Large type of P DLG antibody, LP DLG antibody, P DLG5 antibody, PDLG antibody, Placenta and prostate DLG antibody

UniProt:

Q8TDM6

Application Details

Application Notes:

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Optimal working dilution should be determined by the investigator.