antibodies.com

Datasheet for ABIN7150124 anti-DDB1 antibody (AA 741-943)

2 Images



Overview

Quantity:	100 µg
Target:	DDB1
Binding Specificity:	AA 741-943
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DDB1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human DNA damage-binding protein 1 protein (741-943AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	DDB1
Alternative Name:	DDB1 (DDB1 Products)
Background:	Background: Required for DNA repair. Binds to DDB2 to form the UV-damaged DNA-binding
	protein complex (the UV-DDB complex). The UV-DDB complex may recognize UV-induced DNA

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7150124 | 09/10/2023 | Copyright antibodies-online. All rights reserved. damage and recruit proteins of the nucleotide excision repair pathway (the NER pathway) to initiate DNA repair. The UV-DDB complex preferentially binds to cyclobutane pyrimidine dimers (CPD), 6-4 photoproducts (6-4 PP), apurinic sites and short mismatches. Also appears to function as a component of numerous distinct DCX (DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins. The functional specificity of the DCX E3 ubiquitin-protein ligase complex is determined by the variable substrate recognition component recruited by DDB1. DCX(DDB2) (also known as DDB1-CUL4-ROC1, CUL4-DDB-ROC1 and CUL4-DDB-RBX1) may ubiquitinate histone H2A, histone H3 and histone H4 at sites of UV-induced DNA damage. The ubiquitination of histones may facilitate their removal from the nucleosome and promote subsequent DNA repair. DCX(DDB2) also ubiquitinates XPC, which may enhance DNA-binding by XPC and promote NER. DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitination of TP53 in response to radiation-induced DNA damage and during DNA replication. DCX(ERCC8) (the CSA complex) plays a role in transcription-coupled repair (TCR). May also play a role in ubiquitination of CDKN1B/p27kip when associated with CUL4 and SKP2.

Aliases: Damage specific DNA binding protein 1 antibody, Damage-specific DNA-binding protein 1 antibody, DDB 1 antibody, DDB p127 subunit antibody, Ddb1 antibody, DDB1_HUMAN antibody, DDBa antibody, DNA damage binding protein 1 antibody, DNA damage-binding protein 1 antibody, DNA damage-binding protein a antibody, HBV X-associated protein 1 antibody, UV damaged DNA binding factor antibody, UV damaged DNA binding protein 1 antibody, UV DDB 1 antibody, UV DDB1 antibody, UV-damaged DNA-binding factor antibody, UV-damaged DNAbinding protein 1 antibody, UV-DDB 1 antibody, X associated protein 1 antibody, XAP 1 antibody, XAP-1 antibody, XAP1 antibody, Xeroderma pigmentosum group E complementing protein antibody, XPE BF antibody, XPE binding factor antibody, XPE-BF antibody, XPE-binding factor antibody

UniProt:	
----------	--

Pathways:

DNA Damage Repair

Q16531

Application Details


Application Notes:	Recommended dilution: IHC:1:20-1:200, IF:1:20-1:200,
Restrictions:	For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7150124 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunofluorescence

Image 1. Immunofluorescence staining of Hela cells with ABIN7150124 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Immunohistochemistry

Image 2. Immunohistochemistry of paraffin-embedded human kidney tissue using ABIN7150124 at dilution of 1:100

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7150124 | 09/10/2023 | Copyright antibodies-online. All rights reserved.