

Datasheet for ABIN7150165

**anti-Retinoblastoma Binding Protein 8 antibody (AA 126-331)
(Biotin)**[Go to Product page](#)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µg |
| Target: | Retinoblastoma Binding Protein 8 (RBBP8) |
| Binding Specificity: | AA 126-331 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Retinoblastoma Binding Protein 8 antibody is conjugated to Biotin |
| Application: | ELISA |

Product Details

| | |
|-------------------|--|
| Immunogen: | Recombinant Human DNA endonuclease RBBP8 protein (126-331AA) |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| | |
|-------------------|--|
| Target: | Retinoblastoma Binding Protein 8 (RBBP8) |
| Alternative Name: | RBBP8 (RBBP8 Products) |
| Background: | Background: Endonuclease that cooperates with the MRE11-RAD50-NBN (MRN) complex in processing meiotic and mitotic double-strand breaks (DSBs) by ensuring both resection and |

Target Details

intrachromosomal association of the broken ends. Functions downstream of the MRN complex and ATM, promotes ATR activation and its recruitment to DSBs in the S/G2 phase facilitating the generation of ssDNA. Component of the BRCA1-RBBP8 complex that regulates CHEK1 activation and controls cell cycle G2/M checkpoints on DNA damage. Promotes microhomology-mediated alternative end joining (A-NHEJ) during class-switch recombination and plays an essential role in chromosomal translocations.

Aliases: COM1 antibody, COM1_HUMAN antibody, CtBP interacting protein antibody, CtBP-interacting protein antibody, CtIP antibody, DNA endonuclease RBBP8 antibody, JWDS antibody, RB binding protein 8 endonuclease antibody, RBBP-8 antibody, RBBP8 antibody, Retinoblastoma-binding protein 8 antibody, Retinoblastoma-interacting protein and myosin-like antibody, Rim antibody, SAE2 antibody, SCKL2 antibody, Sporulation in the absence of SPO11 protein 2 homolog antibody

UniProt: [Q99708](#)

Pathways: [Cell Division Cycle](#), [DNA Damage Repair](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.