antibodies -online.com





anti-POLR3K antibody (AA 1-108)



Go to Product page

\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μg
Target:	POLR3K
Binding Specificity:	AA 1-108
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This POLR3K antibody is un-conjugated
Application:	ELISA

Product Details

Immunogen:	Recombinant Human DNA-directed RNA polymerase III subunit RPC10 protein (1-108AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	POLR3K
Alternative Name:	POLR3K (POLR3K Products)
Background:	Background: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA
	using the four ribonucleoside triphosphates as substrates. Component of RNA polymerase III

which synthesizes small RNAs, such as 5S rRNA and tRNAs. Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERs) induce type I interferon and NF- Kappa-B through the RIG-I pathway.

Aliases: C11 antibody, C11 RNP3 antibody, DNA directed RNA polymerase III subunit K antibody, DNA directed RNA polymerase III subunit RPC10 antibody, DNA directed RNA polymerases III 12.5 kDa polypeptide antibody, DNA-directed RNA polymerase III subunit K antibody, DNA-directed RNA polymerase III subunit RPC10 antibody, hRPC11 antibody, HsC11p antibody, My010 antibody, POLR3K antibody, Polymerase (RNA) III (DNA directed) polypeptide K antibody, Polymerase (RNA) III (DNA directed) polypeptide K, 12.3 kDa antibody, RNA polymerase III 12.5 kDa subunit antibody, RNA polymerase III subunit (hRPC11) antibody, RNA polymerase III subunit C10 antibody, RNA polymerase III subunit C11 antibody, RNA polymerase III subunit C11 antibody, RPC11 antibody, RPC12.5 antibody

UniProt:

Q9Y2Y1

Application Details

Application Notes:

Storage Comment:

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C

Ontimal working dilution about the determined by the investigator

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.