antibodies -online.com





anti-POLR3B antibody (AA 831-939)

2 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	POLR3B
Binding Specificity:	AA 831-939
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This POLR3B antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human DNA-directed RNA polymerase III subunit RPC2 protein (831-939AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	POLR3B
Alternative Name:	POLR3B (POLR3B Products)
Background:	Background: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA
	using the four ribonucleoside triphosphates as substrates. Second largest core component of

RNA polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNAs. Proposed to contribute to the polymerase catalytic activity and forms the polymerase active center together with the largest subunit. Pol III is composed of mobile elements and RPC2 is part of the core element with the central large cleft and probably a clamp element that moves to open and close the cleft (By similarity). Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERs) induce type I interferon and NF- Kappa-B through the RIG-I pathway.

Aliases: C128 antibody, DNA directed RNA polymerase III 127.6 kDa polypeptide antibody, DNA directed RNA polymerase III subunit B antibody, DNA-directed RNA polymerase III 127.6 kDa polypeptide antibody, DNA-directed RNA polymerase III subunit B antibody, DNA-directed RNA polymerase III subunit RPC2 antibody, FLJ10388 antibody, polr3b antibody, Polymerase (RNA) III (DNA directed) polypeptide B antibody, RNA polymerase III subunit C2 antibody, RNA polymerase III subunit RPC2 antibody, RPC2 antibody, RPC2_HUMAN antibody


UniProt:


Q9NW08

Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only
Handling	

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.





Immunohistochemistry

Image 1. IHC image of ABIN7150454 diluted at 1:300 and staining in paraffin-embedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunohistochemistry

Image 2. IHC image of ABIN7150454 diluted at 1:300 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.