## antibodies .- online.com





## anti-DAD1 antibody (AA 1-113) (HRP)

DAD1

DAD1 (DAD1 Products)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Target:

Alternative Name:

Background:

Quantity:	100 μg
Target:	DAD1
Binding Specificity:	AA 1-113
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DAD1 antibody is conjugated to HRP
Application:	ELISA
Product Details	
Immunogen:	Recombinant Human Dolichyl-diphosphooligosaccharideprotein glycosyltransferase subunit DAD1 protein (1-113AA)
lmmunogen:	DAD1 protein (1-113AA)
Immunogen: Isotype:	DAD1 protein (1-113AA)  IgG

Background: Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial

transfer of a defined glycan (Glc3Man9GlcNAc2 in eukaryotes) from the lipid carrier dolichol-pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation (PubMed:22467853). N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity (By similarity). Required for the assembly of both SST3A- and SS3B-containing OST complexes. Loss of the DAD1 protein triggers apoptosis (PubMed:22467853).

Aliases: DAD 1 antibody, DAD-1 antibody, dad1 antibody, DAD1\_HUMAN antibody, Defender against cell death 1 antibody, Dolichyl diphosphooligosaccharide protein glycosyltransferase subunit DAD1 antibody, Oligosaccharyl transferase subunit DAD1 antibody, Oligosaccharyl transferase subunit DAD1 antibody, OST2 antibody

UniProt:

P61803

## **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300	
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	