antibodies -online.com





anti-STT3A antibody (AA 39-114)





Overview

| Overview | |
|----------------------|--------------------------------------|
| Quantity: | 100 μg |
| Target: | STT3A |
| Binding Specificity: | AA 39-114 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This STT3A antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC) |
| Product Details | |

| Immunogen: | Recombinant Human Dolichyl-diphosphooligosaccharideprotein glycosyltransferase subunit STT3A protein (39-114AA) |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | STT3A |
|-------------------|---|
| Alternative Name: | STT3A (STT3A Products) |
| Background: | Background: Catalytic subunit of the N-oligosaccharyl transferase (OST) complex which |

catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). SST3A seems to be involved in complex substrate specificity. STT3A is present in the majority of OST complexes and mediates cotranslational N-glycosylation of most sites on target proteins, while STT3B-containing complexes are required for efficient cotranslational glycosylation and mediate glycosylation of sites that have been skipped by STT3A.

Aliases: B5 antibody, Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit STT3A antibody, FLJ27038 antibody, Integral membrane protein 1 antibody, Integral transmembrane protein 1 antibody, ITM1 antibody, MGC9042 antibody, Oligosaccharyl transferase subunit STT3A antibody, STT 3A antibody, STT3 subunit of the oligosaccharyltransferase complex homolog A antibody, STT3-A antibody, STT3A antibody, STT3A_HUMAN antibody, TMC antibody, Transmembrane protein TMC antibody


UniProt:

P46977

Application Details

| Application Notes: | Recommended dilution: IHC:1:20-1:200, |
|--------------------|---------------------------------------|
| Restrictions: | For Research Use only |
| Handling | |
| | |

| Format: | Liquid |
|--------------------|---|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human pancreatic cancer using ABIN7150603 at dilution of 1:100