

Datasheet for ABIN7151144
anti-RNF133 antibody (AA 65-167)



[Go to Product page](#)

3 Images

Overview

| | |
|----------------------|---|
| Quantity: | 100 µg |
| Target: | RNF133 |
| Binding Specificity: | AA 65-167 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RNF133 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF) |

Product Details

| | |
|-------------------|---|
| Immunogen: | Recombinant Human E3 ubiquitin-protein ligase RNF133 protein (65-167AA) |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | Antigen Affinity Purified |

Target Details

| | |
|-------------------|---|
| Target: | RNF133 |
| Alternative Name: | RNF133 (RNF133 Products) |
| Background: | Background: Has E3 ubiquitin-protein ligase activity. Aliases: E3 ubiquitin protein ligase RNF133 antibody, E3 ubiquitin-protein ligase RNF133 |

Target Details

antibody, Goliath-related E3 ubiquitin-protein ligase 2 antibody, Greul2 antibody, MGC27072 antibody, RING finger protein 133 antibody, RN133_HUMAN antibody, RNF 133 antibody, Rnf133 antibody

UniProt: [Q8WVZ7](#)

Application Details

Application Notes: Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of HeLa cells using ABIN7151144 at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western Blotting

Image 2. Western blot All lanes: RNF133 antibody at 2 μ g/mL + K562 whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 43 kDa Observed band size: 43 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human testis tissue using ABIN7151144 at dilution of 1:100