

Datasheet for ABIN7151354
anti-ART5 antibody (AA 45-291)[Go to Product page](#)

4 Images

Overview

Quantity:	100 µg
Target:	ART5
Binding Specificity:	AA 45-291
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ART5 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Ecto-ADP-ribosyltransferase 5 protein (45-291AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	ART5
Alternative Name:	ART5 (ART5 Products)
Background:	ADP ribosyltransferase 5 antibody, ADP-ribosyltransferase C2 and C3 toxin-like 5 antibody, ART 5 antibody, ART5 antibody, ARTC5 antibody, EC 2.4.2.31 antibody, Ecto ADP ribosyltransferase

Target Details

5 antibody, Ecto-ADP-ribosyltransferase 5 antibody, MGC22848 antibody, Mono (ADP ribosyl) transferase 5 antibody, Mono(ADP-ribosyl)transferase 5 antibody, NAD(P)(+) arginine ADP ribosyltransferase 5 antibody, NAD(P)(+)-arginine ADP-ribosyltransferase 5 antibody, NAR5_HUMAN antibody

UniProt: [Q96L15](#)

Application Details

Application Notes: Recommended dilution: IHC:1:100-1:1000,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human brain tissue using ABIN7151354 at dilution of 1:100



Immunohistochemistry

Image 2. Immunohistochemistry of paraffin-embedded human heart tissue using ABIN7151354 at dilution of 1:100



Immunohistochemistry

Image 3. IHC image of ABIN7151354 diluted at 1:1000 and staining in paraffin-embedded human pancreatic cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN7151354.