antibodies -online.com





anti-PRODH antibody (AA 141-240) (Biotin)



Overview

Quantity:	100 μL
Target:	PRODH
Binding Specificity:	AA 141-240
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRODH antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PRODH
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Horse
Purification:	Purified by Protein A.

Target Details

Target:	PRODH
Alternative Name:	PRODH (PRODH Products)

Target Details

Background:	Synonyms: POX, PIG6, HSPOX2, PRODH1, PRODH2, TP53I6, Proline dehydrogenase 1,
	mitochondrial, Proline oxidase, Proline oxidase 2, p53-induced gene 6 protein, PRODH, POX2
	Background: Proline oxidase catalyzes the conversion of proline to pyrroline-5-carboxylate, or
	P5C during the degradation of the amino acid Proline. Defects in PRODH are the cause of
	hyperprolinemia type 1, a disorder characterized by elevated serum proline levels. Defective
	PRODH may be involved in the psychiatric and behavioral phenotypes associated with the
	22q11 velocardiofacial and DiGeorge syndrome and may be associated with susceptibility to
	schizophrenia 4 (SCZD4).
Gene ID:	5625
UniProt:	043272
Pathways:	Response to Water Deprivation, Monocarboxylic Acid Catabolic Process
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months